



## The Setting

Riverside County is located within three air basins, as can be seen on Figure AQ-1, Riverside County Air Quality Basins. They are the South Coast Air Basin (SOCAB), Salton Sea Air Basin (SSAB) and the Mojave Desert Air Basin (MDAB). Air quality within each basin is not only affected by various emissions sources (mobile, industry, etc.), but also by atmospheric conditions such as wind speed, wind direction, temperature and rainfall. The following provides a description of each air basin and its relevant climate and meteorological conditions affecting air pollution.

### SOUTH COAST AIR BASIN

Western Riverside County (west of the San Geronio Pass) is located within the South Coast Air Basin (SOCAB), which includes all of Orange County and the non-desert portions of Los Angeles, Riverside and San Bernardino Counties. Air quality conditions in the SOCAB are under the jurisdiction of the South Coast Air Quality Management District (SCAQMD).

According to the Air Quality Management Plan (AQMP), the worst air quality problem in the nation occurs in the South Coast Air Basin. With very light average wind speeds, the basin atmosphere has a limited capability to disperse air contaminants horizontally. The dominant daily wind pattern is a daytime sea breeze (onshore breeze) and a nighttime land breeze (offshore breeze), broken only occasionally by winter storms and infrequent strong Santa Ana winds from the Great Basin, Mojave, and deserts to the north.

On virtually all spring and early summer days, most of the pollution produced during an individual day is moved out of the basin through mountain passes, or is lifted by the warm, vertical currents produced by the heating of mountain slopes. In those seasons, the basin can be “flushed” of pollutants by a transport of ocean air during the afternoon. From late summer through the winter months, the flushing is less pronounced because of lower wind speeds and the earlier appearance of offshore winds. With extremely stagnant wind flows, the drainage winds may begin near the mountains by late afternoon. Remaining pollutants are trapped and begin to accumulate during the night and the following morning. A low average morning wind speed in pollution source areas is an important indicator of air stagnation potential.

The vertical dispersion of air pollutants in the South Coast Air Basin is hampered by the presence of a temperature inversion in the layers of the atmosphere near the surface of the Earth. In a normal situation, as temperatures decrease with altitude, air continues to rise as it remains warmer than the surrounding air. With an inversion layer, air cannot continue to expand upwards, as it is trapped by the warmer air above.



**Santa Ana Winds** - Santa Ana winds are generally defined as warm, dry winds that blow from the east or northeast (offshore) occurring predominantly between the months of December and February. The winds develop when a region of high pressure builds over the Great Basin (the high plateau east of the Sierra Mountains and west of the Rocky Mountains including most of Nevada and Utah) and move locally across the Mojave Desert and then over and through passes in the San Gabriel, San Bernardino and San Jacinto Mountains.



**Inversion layer** - A layer of warm air that traps the cooler air and any pollutants it carries below.

However, as the day progresses and the sun warms the ground, the surface layer of air approaches a temperature equal to that of the inversion layer. When these temperatures become equal, the inversion layer begins to erode at its lower edge. If enough warming takes place, the inversion layer becomes weaker and weaker and finally “breaks.” The surface air layers can then mix upward without limit. This phenomenon is frequently observed in the middle of late afternoon on hot summer days when the smog appears to clear up suddenly. Winter inversions frequently break by mid-morning, thereby preventing contaminant build-up.

The combination of low wind speeds and low level inversions produces the greatest concentration of pollutants. On high wind days other air pollutants including particulate matter such as dust and soil are swept and carried in the air. On days of no inversion or on days of winds averaging over 15 miles per hour, there will be no important smog effects, during either summer or winter.



**Smog** - A combination of smoke, ozone, hydrocarbons, nitrogen oxides, and other chemically reactive compounds which, under certain conditions of weather and sunlight, may result in a murky brown haze that causes adverse health effects. The primary source of smog in California is motor vehicles.

In the winter, the greatest pollution problems are carbon monoxide and oxides of nitrogen because of extremely low level inversions and air stagnation during the night and early morning hours. Smog levels are much lower during this season due to the lack of strong inversion during the daylight hours and the lack of intense sunlight which is needed to produce photochemical reactions.

In the summer, the longer daylight hours and the brighter sunshine combine to cause a reaction between hydrocarbons and oxides of nitrogen to form more smog. Carbon monoxide is not as great a problem in summer because inversions are not as low and intense in the surface boundary layer (within 100 feet of the ground) as in winter and because horizontal ventilation is better in summer.

The basin-wide average occurrence of inversion at the ground surface is 11 days per month; the averages vary from two days in June to 22 days in December and January. The potential for high concentration varies seasonally for many contaminants. During late spring, summer and early fall, light winds, low mixing heights and brilliant sunshine combine to produce conditions favorable for the maximum production of photochemical oxidants, mainly ozone. During the spring and summer, when fairly deep marine layers are frequently found in the Basin, sulfate concentrations are at their peak.

## SALTON SEA AIR BASIN



**Subtropical High Cell** - An area of atmospheric high pressure located at approximately 30 degrees north and south latitude. Air tends to sink near high-pressure centers, which inhibits precipitation and cloud formation. This is why high-pressure systems tend to bring bright, sunny days with calm weather.

The middle part of Riverside County (between San Geronio Pass and Joshua Tree National Monument), belongs in the Salton Sea Air Basin (SSAB), along with Imperial County. Air quality conditions in this portion of the County, although in the SSAB, are also administered by the SCAQMD. The SCAQMD is responsible for the development of the regional Air Quality Management Plan and efforts to regulate pollutant emissions from a variety of sources.

The SSAB portion of Riverside County is separated from the SOCAB region by the San Jacinto Mountains and from the Mojave Desert Air Basin to the east by the Little San Bernardino Mountains. During the summer, the SSAB is generally influenced by a Pacific Subtropical High Cell that sits off the coast, inhibiting cloud formation and encouraging daytime solar heating. The SSAB is rarely influenced by cold air masses moving south from Canada and Alaska, as these systems are weak and diffuse by the time they reach the desert. Most desert